

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## **IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
Please do not report the images to the  
Image Problem Mailbox.

**THIS PAGE BLANK (USPTO)**



Europäisches  
Patentamt

Eur pean  
Patent Office

Office eur péen  
des brevets

REC'D 03 JUL 2000

EPO - DG 1

WIPO

PCT 02.06.2000

(70)

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99111321.8

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE,  
LA HAYE, LE

12/04/00





Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.: 99111321.8  
Demande n°:

Anmeldetag:  
Date of filing: 10/06/99  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
Stymne, Sten

268 90 Svalöv

SWEDEN

Dahlqvist, Anders

244 66 Furulund

SWEDEN

Bezeichnung der Erfindung:

Title of the invention:

Titre de l'invention:

A new class of enzymes in the biosynthetic pathway for the production of triacylglycerol and recombinant DNA molecules encoding these enzymes

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat: EP  
State:  
Pays:

Tag: 01/04/99  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

EPA 99106656

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

/

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

Further applicants are : Ulf STAHL, 753 24 UPPSALA, Sweden  
Marit LENMAN, 233 59 LUND, Sweden  
Antoni BANAS, 08-110 SIEDLCE, Poland  
Hans RONNE, 756 54 UPPSALA, Sweden



A new class of enzymes in the biosynthetic pathway for the production of triacylglycerol and recombinant DNA molecules encoding these enzymes

Applicants and inventors:

Anders Dahlqvist, Ulf Ståhl, Marit Lenman, Antoni Banas, Hans Ronne and Sten Stymne

Description

The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding enzymes catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

The invention further relates to novel type of enzymes, the genes encoding such enzymes and the use of these genes for transformation. More specifically, the invention relates to use of a previously un-described type of enzymes hereinafter designated phospholipid:diacylglycerol acyltransferases (PDAT) and the genes encoding them. This type of genes expressed alone in transgenic organisms will enhance the total amount of oil (triacylglycerols) produced in the cells. The PDAT genes, in combination with a gene for the synthesis of an uncommon fatty acid will, when expressed in transgenic organisms, enhance the levels of the uncommon fatty acids in the triacylglycerols.

There is considerable interest world-wide in producing chemical feedstock, such as fatty acids, for industrial use from renewable plant resources rather than non-renewable petrochemicals. This concept has broad appeal to manufacturers and consumers on the basis of resource conservation and provides significant opportunity to develop new industrial crops for agriculture.

There is a diverse array of unusual fatty acids in oils from wild plant species and these have been well characterised (see e.g. Badami & Patil, 1981). Many of these acids have industrial potential and this has led to interest in domesticating relevant plant species to enable agricultural production of particular fatty acids.

Development in genetic engineering technologies combined with greater understanding of the biosynthesis of unusual fatty acids now makes it possible to transfer genes coding for key enzymes involved in the synthesis of a particular fatty acid from a wild species into domesticated oilseed crops. In this way individual fatty acids can be produced in high purity and quantities at moderate costs.

In all crops like rape, sunflower, oilpalm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds like starch, protein, and fibre is regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of crop. If genes, regulating the allocation of reduced carbon into the production of oil can be up-regulated, the cells will accumulate more oil on the expense of other products. Such genes might not only be used in already high oil producing cells such as oil crops but could also induce significant oil production in moderate or low oil containing crops such as e.g. soy, oat, maize, potato, sugarbeets, and turnips as well as in micro-organisms.

#### Summary of the invention

Many of the uncommon fatty acids of interest, e.g. medium chain fatty acids, hydroxy fatty acids, epoxy fatty acids and acetylenic fatty acids, have physical properties that are distinctly different from the common plant fatty acids. The present inventors have found that, in plant species naturally accumulating these uncommon fatty acids in their seed oil (i.e. triacylglycerol), these acids are absent, or present in very low amounts in the membrane (phospho)lipids of the seed. The low concentration of these acids in the membrane lipids is most likely a prerequisite for proper membrane function and thereby for proper cell functions. One aspect of the invention is that seeds of transgenic crops can be made to accumulate high amounts of uncommon fatty acids if these fatty acids are efficiently removed from the membrane lipids and channelled into seed triacylglycerols.

The inventors have identified a novel class of enzymes in plants catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the production of triacylglycerol and, presumably, lysophospholipids and that these enzymes (phospholipid:diacylglycerol acyltransferases abbreviated as PDAT) are involved in the removal of hydroxylated, epoxygenated fatty acids, and probably also other uncommon fatty acids such as medium chain fatty acids, from phospholipids in plants. Further, the same enzyme reaction was shown to be present in microsomal preparations from baker's yeast (*Saccharomyces cerevisiae*). A so called 'knock out' yeast mutant, disrupted in the respective gene was obtained and microsomal membranes from the mutant was shown to totally lack PDAT activity. Thus, it was proved that the disrupted gene encodes a PDAT enzyme (SEQ ID NO: 1 and 2). In addition, a gene from *Schizosaccharomyces pombe*, SPBC776.14 (SEQ ID. NO. 3), having 46% identity over 571 amino acids, and three *Arabidopsis thaliana* genomic sequences coding for putative proteins with amino acid sequence having approximately 44 % identity over a total of 378 amino acids (SEQ ID NO: 4), 47 % identity over 73 amino acids (SEQ ID NO: 9), and 33% identity over 105 amino acids (SEQ ID NO. 10), with the yeast enzyme, were



identified.

Also, a partially sequenced cDNA clone from *Neurospora crassa* (SEQ ID NO. 8) showed 42% identity over 167 amino acids, and a *Zea mays* est clone (SEQ ID NO. 6 and 7) showed 36% identity over 96 amino acids, with the yeast enzyme were identified. Finally, two cDNA clones were identified, one *Arabidopsis thaliana* est (SEQ ID NO. 5) which was 98% identical in nucleotide sequence to the *Arabidopsis thaliana* genomic sequence (ID no. 4), and a *Lycopersicon esculentum* est clone (SEQ ID NO. 11) showed 84% identity in 72 nucleotides to Seq ID No. 10.

In a first embodiment, this invention is directed to a PDAT enzyme and nucleic acid sequences that encode a PDAT. This includes biologically active PDATs and encoding sequences for PDATs as well as sequences that are to be used as probes, vectors for transformation or cloning intermediates. The PDAT encoding sequence may encode a complete or partial sequence depending upon the intended use. All or a portion of the genomic sequence, cDNA sequence, precursor PDAT or mature PDAT is intended.

In a different aspect, this invention relates to a method for producing a PDAT in a host cell or progeny thereof, including genetically engineered oil seeds, yeast and moulds or any other oil accumulating organism, via the expression of a construct in the cell. Cells containing a PDAT as a result of the production of the PDAT encoding sequence are also contemplated within the scope of the invention.

In a different embodiment, this invention also relates to methods of using a DNA sequence encoding a PDAT for increasing the oil-content within a cell.

Another aspect of the invention relates to the accommodation of high amounts of uncommon fatty acids in the triacylglycerol produced within a cell, by introducing a DNA sequence producing a PDAT that specifically removes these fatty acids from the membrane lipids of the cell and channel them into triacylglycerol. Plant cells having such a modification are also contemplated herein.

A PDAT of this invention includes any sequence of amino acids, such as a protein, polypeptide or peptide fragment obtainable from a microorganism, animal or plant source that demonstrates the ability to catalyse the production of triacylglycerol from a phospholipid and diacylglycerol under enzyme reactive conditions. By "enzyme reactive conditions" is meant that any necessary conditions are available in an environment (e.g., such factors as temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

Other PDATs are obtainable from the specific sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic PDATs, including modified amino acid sequences and starting materials for synthetic-protein modelling from the exemplified PDATs and from PDATs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated, increased and the like, whether such sequences were partially or wholly synthesised. Sequences that are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method used to obtain the protein or sequence, are equally considered naturally derived.

The nucleic acid sequences (DNA and RNA) of the present invention can be used to screen and recover "homologous" or "related" PDATs from a variety of plant and microbial sources.

Further, it is also apparent that a person skilled in the art can, with the information provided in this application, in any organism identify a PDAT activity, purify an enzyme with this activity and thereby identify a "non-homologues" nucleic acid sequence encoding such an enzyme.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1.

PDAT activity in microsomal fractions of *S. cerevisiae*. Aliquots of lyophilised microsomal membranes (10 nmol phosphatidylcholine) from a wild type yeast (strain YN979) (lane 1-3, 3'), a yeast mutant (strain B10280), disrupted in the YNRO08w gene (lane 4-6, 9) or the yeast mutant complemented with a single copy plasmid containing the PDAT gene (lane 7) were assayed for PDAT activity. 2 nmol sn-1-oleoyl-sn-2-[<sup>14</sup>C]-ricinoleoylphosphatidylcholine (lane 1-7) or sn-1-oleoyl-sn-2-[<sup>14</sup>C]-oleoylphosphatidylcholine (lane 8-9) and 5 nmol of dioleoyl-diacylglycerol (lane 2, 5, 7-9) or rac-oleoyl-vernoleoyldiacylglycerol (lane 3, 6) were added in benzene solution. The benzene was evaporated under N<sub>2</sub> (g) and 0.1 ml of 50 mM potassium phosphate, pH 7.2, was added. The suspension was thoroughly mixed and after 90 min at 30 °C the lipids were extracted in chloroform and separated on thin layer chromatography on silica gel 60 plates in hexan/diethylether/acetic acid (35:70:1.5). The radioactive lipids were visualised and quantified on the plates by electronic autoradiography (Instant imager, Packard, US). Abbreviations used: triacylglycerol, TAG, FA, fatty acid (i.e. oleic acid); 1-OH-TAG, monoricinoleoyl-triacylglycerol; 1-OH-1-epTAG, monoricinoleoyl-monovernoleoyl-triacylglycerol and OH-FA, ricinoleic acid.

Brief Description of the SEQ ID:

SEQ ID NO. 1: Genomic DNA sequence of the *Saccharomyces cerevisiae* PDAT gene, YNR008w, genebank nucleotide ID number 1302481, and the suggested YNR008w amino acid sequence.

SEQ ID NO. 2: The suggested amino acid sequence of the yeast gene YNR008w from *Saccharomyces cerevisiae*.

SEQ ID NO. 3: Genomic DNA sequence of the *Schizosaccharomyces pombe* gene SPBC776.14.

SEQ ID NO. 4: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AB006704.

SEQ ID NO. 5: Nucleotide sequence and the corresponding amino acid sequence of the *Arabidopsis thaliana* est clone with genebank accession number T04806, and ID number 315966.

SEQ ID NO. 6: Nucleotide and amino acid sequence of the *Zea mays* cDNA clone with genebank ID number g4388167.

SEQ ID NO. 7: Amino acid sequence of the *Zea mays* cDNA clone with genebank ID number g4388167.

SEQ ID NO. 8: DNA sequence of part of the *Neurospora crassa* cDNA clone W07G1, ID number g4241729.

SEQ ID NO. 9: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC004557.

SEQ ID NO. 10: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC003027.

SEQ ID NO. 11: DNA sequence of part of the *Lycopersicon esculentum* cDNA clone with genebank accession number AI486635.

The present invention can be essentially characterized by the following aspects:

1. Use of a PDAT PDAT gene (genomic clone or cDNA) for transformation.
2. Use of a DNA molecule according to item 1 wherein said DNA is used for transformation of any organism in order to be expressed in this organism and result in an active recombinant PDAT enzyme in order to increase oil content of the organism.
3. Use of a DNA molecule of item 1 wherein said DNA is used for transformation of any organism in order to prevent the accumulation of undesirable fatty acids in the membrane lipids.

4. Use according to item 1, wherein said PDAT gene is used for transforming transgenic oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, such as medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
5. Use according to item 1, wherein said PDAT gene is used for transforming organisms, and wherein said organisms are crossed with other oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, comprising medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
6. Use according to item 1, wherein the enzyme encoded by said PDAT gene or cDNA is coding for a PDAT with distinct acyl specificity.
7. Use according to item 1 wherein said PDAT encoding gene or cDNA, is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 30% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
8. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
9. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 60% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
10. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 80% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
11. Use according to claim 1 wherein said PDAT encoding gene or cDNA is derived from plants or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence encoded by the PDAT gene from *Arabidopsis thaliana* as presented in SEQ. ID. NO. 4, 9 or 10, or to the protein encoded by the fulllength counterpart of the partial *Zea mays*, *Lycopersicon esculentum*, or *Neurospora crassa* cDNA clones.
12. Transgenic oil accumulating organisms comprising, in their genome, a PDAT gene transferred by recombinant DNA technology or somatic hybridization.
13. Transgenic oil accumulating organisms according to item 12 comprising, in their genome, a PDAT gene having specificity for substrates with particular uncommon fatty acid and the gene for said uncommon fatty acid.
14. Transgenic organisms according to item 12 or 13 which are selected from the group consisting of fungi, plants and animals.

15. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants.
16. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a storage organ specific promotor.
17. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a seed promotor.
18. Oils from organisms according to item 12 - 17.
19. A method for altering acyl specificity of a PDAT by alteration of the nucleotide sequence of a naturally occurring encoding gene and as a consequence of this alternation creating a gene encoding for an enzyme with novel acyl specificity.
20. A protein encoded by a DNA molecule according to item 1 or a functional fragment thereof.
21. A protein of item 20 designated phospholipid:diacylglycerol acyltransferase.
22. A protein of item 21 which has a distinct acyl specificity.
23. A protein of item 13 having the amino acid sequence as set forth in Seq id no. 2, (and the proteins encoded by the genes fulllength or partial genes set forth in SEQ ID NO. 1, 3, 4, 5, 6, 7, 8, 9, 10 or 11) or an amino acid sequence with at least 30 % homology to said amino acid sequence.
24. A protein of item 23 isolated from *Saccharomyces cerevisiae*.

#### EXAMPLE 1

Determination of substrate utilisation by the acyl-CoA independent acyltransferase in the synthesis of triacylglycerols in microsomal preparations of developing castor bean endosperm and bakers yeast.

Microsomal membranes prepared from developing endosperm of castor bean (*Ricinus communis*) catalyse the selective transfer of ricinoleoyl-(12-hydroxy-9-octadecenoyl) and vernoloyl(12-epoxy-9-octadecenoyl) groups from both diacylglycerols and phosphatidylcholine into triacylglycerols. The substrate utilisation was investigated in castor bean microsomes by using radioactive sn-1-oleoyl-sn-2-<sup>14</sup>C]ricinoleoyl-diacylglycerol (sn-2-<sup>14</sup>C]-ricinoleoyl-diacylglycerol) or sn-1-oleoyl-sn-2-<sup>14</sup>C]ricinoleoyl-phosphatidylcholine (sn-2-<sup>14</sup>C]-ricinoleoyl-phosphatidylcholine) together with different non-radioactive diacylglycerol (see Table 1).

The preparatian of microsomal fractions of developing castor bean endosperm and freeze drying of the microsomes were performed in known manner. Assays with addition of diacylglycerol and phosphatidylcholine substrates were performed. The results is presented in Table 1 and showed that if radioactive sn-2-<sup>14</sup>C]ricinoleoyl-diacylglycerol was used as the only added

substrate, 2.8 % of the radio-labelled ricinoleoyl chains were found in triacylglycerol with one ricinoleoyl group, 12.4 % of the radioactivity was found in triacylglycerol-species with two ricinoleoyl groups and only trace amounts were associated with triacylglycerol consisting of three ricinoleoyl groups. If incubations with sn-2-[<sup>14</sup>C] ricinoleoyl-diacylglycerol were performed in a 1:4 (mol:mol) mixture with non-radioactive diacylglycerol species containing one vernoloyl group, the distribution of radioactivity between different molecular species of triacylglycerol changed only marginally compared to incubations with just radioactive substrate. Only 1.3% of the added <sup>14</sup>C-labelled ricinoleoyl groups were metabolised into triacylglycerol species with one ricinoleoyl and one vernoloyl group. Similarly, only marginal changes in the radioactive triacylglycerol molecular species was seen in incubations where sn-2-[<sup>14</sup>C]-ricinoleoyl-diacylglycerol was mixed with non-labelled divernoloyl-diacylglycerol. However, by adding unlabelled diricinoleoyl-phosphatidylcholine together with sn-2-[<sup>14</sup>C]-ricinoleoyl-diacylglycerol the radioactivity metabolised into the different triacylglycerol species were substantially altered.

Only trace amounts of radioactivity were detected in triacylglycerol species with one ricinoleoyl chain whereas the radioactivity in triacylglycerol with two ricinoleoyl groups were doubled as compared to incubations with only sn-2-[<sup>14</sup>C]-ricinoleoyl-diacylglycerol added.

Table 1. *In vitro* synthesis of triacylglycerols in microsomal preparations of developing castor bean. Aliquots of microsomes (20 nmol PC) were lyophilised and substrate lipids were added in benzene solution: (A) 0.4 nmol [ $^{14}$ C]-DAG (7760 dpm/nmol) and where indicated 1.6 nmol unlabelled DAG; (B) 0.4 nmol [ $^{14}$ C]-DAG (7760 dpm/nmol) and 5 nmol unlabelled di-ricinoleoyl-PC and (C) 0.25 nmol [ $^{14}$ C]-PC (4000 dpm/nmol) and 5 nmol unlabelled DAG. The benzene was evaporated by  $N_2$  and 0.1 ml of 50 mM potassium phosphate was added, thoroughly mixed and incubated at 30 °C for (A) 20 min; (B) and (C) 30 min. Assays were terminated by extraction of the lipids in chloroform (Dahlqvist et al. 1998). The lipids were then separated by thin layer chromatography on silica gel 60 plates (Merck, Darmstadt, Germany) in hexan/diethylether/acetic acid 35:70:1.5. The radioactive lipids were visualised and the radioactivity quantified on the plate by electronic autoradiography (Instant Imager, Packard, US). Results are presented as mean values of two experiments.

Substrate added [ $^{14}$ C]-lipid <sup>(a)</sup>	unlabelled lipid <sup>(a)</sup>	mol% of added [ $^{14}$ C]-acyl group in TAG <sup>(b)</sup>				
		1-OH-TAG	2-OH-TAG	1-OH-1-ver-TAG	1-OH-2-ver-TAG	3-OH-TAG
A	mono-[ $^{14}$ C]-ricinoleoyl-DAG	2.8	12.4	-	-	-
	mono-[ $^{14}$ C]-ricinoleoyl-DAG	3.2	12.1	1.3	-	-
	mono-[ $^{14}$ C]-ricinoleoyl-DAG	4	10	0.5	1.2	-
B	mono-[ $^{14}$ C]-ricinoleoyl-DAG	0.3	24.8	-	-	-
	none	6.8	8.0	-	-	4.7
C	mono-[ $^{14}$ C]-ricinoleoyl-PC	8.6	9.8	-	-	5.0
	mono-[ $^{14}$ C]-ricinoleoyl-PC	5.7	16.7	-	-	1.9
	mono-[ $^{14}$ C]-ricinoleoyl-PC	4.5	9.4	-	-	9.5
	mono-[ $^{14}$ C]-ricinoleoyl-PC	6.0	11.5	10.9	0.5	7.4
	mono-[ $^{14}$ C]-ricinoleoyl-PC	6.7	10.8	1.1	8.4	6.8

Radioactivity in different triacylglycerols (TAG) species formed. Abbreviations used: 1-OH-, mono-ricinoleoyl-, 2-OH-, di-ricinoleoyl-, 3-OH-, tri-ricinoleoyl; 1-OH-1-ver-, mono-ricinoleoyl-monovermoleoyl-, 1-OH-2-ver-, mono-ricinoleoyl-divermoleoyl-. Radiolabelled DAG and PC were prepared enzymatically. The radiolabelled ricinoleoyl group is attached at the *sn*-2-position of the lipid and unlabelled oleoyl group at the *sn*-1-position. Unlabelled DAG with vermoleoyl- or ricinoleoyl chains were prepared by the action of TAG lipase (REF) on oil of *Euphorbia lagascae* or Castor bean, respectively. Synthetic di-ricinoleoyl-PC was kindly provided from Metapontum Agribios (Italy)

## EXAMPLE 2

### Transformation and expression of YNROORw gene in yeast

The yeast mutant (strain B 10280) disrupted in the YNRO08w gene, was transformed with the single copy plasmid pFL39 having the PDAT-gene (YNRO08w) under the control of the endogenous promotor region (583 bp 5' untranslated) inserted into the cloning cassette. The transformed yeast was pre-cultivated at 28 °C for 20 h in defined YNB medium without tryptophane added. Cells were harvested and re-suspended in minimal medium (Meesters et al., 1996), supplemented with 16 g/l glycerol to the original volume of the growth culture. The culture was further incubated for 24 h after which cells were harvested by centrifugation. Microsomal fraction of the yeast was prepared as described in Example 1 above and was incubated in the presence of sn-2-[<sup>14</sup>C]-ricinoleoyl-phosphatidylcholine (Fig 1, lane 7). This experiment clearly shows that the PDAT activity could be restored by the expression of the YNRO08w gene in the mutant yeast strain B10280 normally lacking the PDAT-activity.

The effect of the over-expression of the PDAT gene on the lipid accumulation was studied by transforming the wild-type yeast (strain SCY62) with a plasmid pJN92 containing the PDAT gene (YNRO08w) under the control of a GAL1-promotor. The transformed yeast was then cultivated at 28 °C in defined YNB medium lacking uracil. The expression of the PDAT gene was induced by the addition of 2 % (v/v) galactose after 10 hours growth and was further incubated for 18 hours. The yeast cells were harvested and the lipid content of the yeast was analysed by thin layer chromatography and gas liquid chromatography. The results are presented in Table 2. The total lipid content in the yeast with the over-expressed PDAT was 1.3 fold higher than in the control yeast transformed with an empty plasmid pJN92. The expression of the PDAT gene had no effect on the growth rate as determined by optical density measurements. The elevated lipid content in the yeast transformed with PDAT as compared to the control yeast can be totally accounted for by an 80 % increase seen in the triacylglycerol content. The levels of the polar lipids and sterol esters were not significantly effected by the over-expression of the PDAT gene. Hence, these results clearly demonstrate the use of the PDAT gene in increasing the oil content in transgenic organisms.



**Table 2. Lipid content in yeast with overexpressed PDAT gene.**

Yeast transformed with the PDAT gene under the control of a gal promotor in a xxxx plasmid were grown in YNB medium at 28 °C. After after 10 hours growth the expression of the PDAT gene was induced by the addition of 2 % (v/v) galactose. Cells were harvested after additional 18 hours growth and analysed for its lipid content. Yeast transformed with an empty plasmid grown under identical conditions was used as control. Lipids were extracted in chloroform, fractionated on TLC and quantified by GC analyses. The lipid content was measured as nmol fatty acids (FA) per mg yeast (dry weight).

	PDAT-transformed yeast		Control yeast	
	nmol FA/mg	(%)	nmol FA/mg	(%)
Polar lipids	60	32	60	42
Sterol esters	17	9	20	14
Triacylglycerol	105	56	58	40
Other lipids	6	3	6	4
Sum	188		144	

Claims

1. An enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol forming triacylglycerol.
2. A recombinant DNA molecule encoding the enzyme of claim 1.
3. A recombinant DNA molecule according to claim 2 encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, wherein said enzyme comprises an amino acid sequence set forth in SEQ ID NO. 2, or fragments thereof, or is encoded by the genes set forth in SEQ ID NO. 1, 3, 4, 9 or 10, or is encoded by the fulllength genes corresponding to the partial nucleotide sequences set forth in SEQ ID NO. 5, 6, 7, 8, and 11 or sequences 40% or more identical to said sequences.
4. The DNA molecule of claim 2 or 3, wherein said enzyme is designated as phospholipid:diacylglycerol acyltransferase.
5. A vector comprising a DNA molecule of claim 2 or 3.
6. A vector of claim 5 further comprising a selectable marker gene.
7. A host cell containing a DNA molecule of claim 2 or 3.
8. The host cell of claim 7 which is a plant cell or yeast cell.
9. A process for the production of transgenic yeast cells, plant cells or plants comprising a) transforming a DNA molecule of claims 2 or 3 into plant cells or plants; and b) selecting of transformed plant cells or plants having an altered biosynthetic pathway in the production of triacylglycerol.
10. A process of claim 9 wherein the altered biosynthetic pathway is characterised by an increased or altered oil content.
11. A process of claim 9 wherein the altered biosynthetic pathway is characterised by the prevention of accumulation of undesirable fatty acids in the membrane lipids.

12. A protein encoded by a DNA molecule according to claim 2 or 3 or a functional fragment thereof.
13. A method for production of triacylglycerol, comprising enzymatic catalysation by an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol forming triacylglycerol.
14. A method according to claim 13, comprising enzymatic catalysation by a protein or functional fragment according to claim 12.

The present invention relates to a novel class of enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol. Further, the invention relates to the isolation, identification, characterization and use of recombinant DNA molecules encoding this class of enzymes.

## SEQUENCE LISTING

&lt;110&gt; Stymne Dr., Sten

<120> RECOMBINANT DNA MOLECULES ENCODING A NEW CLASS OF  
ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION  
OF TRIACYLGLYCEROL

&lt;130&gt; stymne

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 11

&lt;170&gt;

&lt;210&gt; 1

&lt;211&gt; 1986

&lt;212&gt; genomic DNA

&lt;213&gt; Saccharomyces cerevisiae

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1983)

&lt;400&gt; 1

atg	ggc	aca	ctg	ttt	cga	aga	aat	gtc	cag	aac	caa	aag	agt	gat	tct	48
Met	Gly	Thr	Leu	Phe	Arg	Arg	Asn	Val	Gln	Asn	Gln	Lys	Ser	Asp	Ser	
1					5				10						15	

gat	gaa	aac	aat	aaa	ggg	ggt	tct	gtt	cat	aac	aag	cga	gag	agc	aga	96
Asp	Glu	Asn	Asn	Lys	Gly	Gly	Ser	Val	His	Asn	Lys	Arg	Glu	Ser	Arg	
				20				25					30			

aac	cac	att	cat	cat	caa	cag	gga	tta	ggc	cat	aag	aga	aga	agg	ggt	144
Asn	His	Ile	His	His	Gln	Gln	Gly	Leu	Gly	His	Lys	Arg	Arg	Arg	Gly	
				35				40					45			

att	agt	ggc	agt	gca	aaa	aga	aat	gag	cgt	ggc	aaa	gat	ttc	gac	agg	192
Ile	Ser	Gly	Ser	Ala	Lys	Arg	Asn	Glu	Arg	Gly	Lys	Asp	Phe	Asp	Arg	
				50				55				60				

aaa	aga	gac	ggg	aac	ggt	aga	aaa	cgt	tgg	aga	gat	tcc	aga	aga	ctg	240
Lys	Arg	Asp	Gly	Asn	Gly	Arg	Lys	Arg	Trp	Arg	Asp	Ser	Arg	Arg	Leu	
				65				70				75			80	

att	ttc	att	ctt	ggt	gca	ttc	tta	ggt	gta	ctt	ttg	ccg	ttt	agc	ttt	288
Ile	Phe	Ile	Leu	Gly	Ala	Phe	Leu	Gly	Val	Leu	Leu	Pro	Phe	Ser	Phe	
				85				90						95		

ggc	gct	tat	cat	gtt	cat	aat	agc	gat	agc	gac	ttg	ttt	gac	aac	ttt	336
Gly	Ala	Tyr	His	Val	His	Asn	Ser	Asp	Ser	Asp	L	u	Phe	Asp	Asn	Phe
				100				105					110			

gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt	384
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val	
115 120 125	
ctc cca caa ggt ata agt tgc ttt att gat gat att cag gct ggt aac	432
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn	
130 135 140	
tac tcc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt	480
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly	
145 150 155 160	
aaa caa ctc tta cgt gat tat aat atc gag gcc aaa cat cct gtt gta	528
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val	
165 170 175	
atg gtt cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att	576
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile	
180 185 190	
gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg	624
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp	
195 200 205	
gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg	672
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp	
210 215 220	
ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac	720
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn	
225 230 235 240	
ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat tat ttc atc	768
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile	
245 250 255	
gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att	816
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile	
260 265 270	
ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt	864
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu	
275 280 285	

gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag 912  
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys  
 290 295 300

gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta 960  
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu  
 305 310 315 320

att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg 1008  
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp  
 325 330 335

gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt 1056  
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
 340 345 350

aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc 1104  
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
 355 360 365

gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc 1152  
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
 370 375 380

att caa tta aat acg tta gcc atg tat ggt ttg gaa aag ttc ttc tca 1200  
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
 385 390 395 400

aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca 1248  
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
 405 410 415

atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct 1296  
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
 420 425 430

tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att 1344  
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
 435 440 445

cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg 1392  
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
 450 455 460

aaa gac gcc att aac atg aca tta tgc ata tca cct gaa tgg ctc caa 1440  
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
 465 470 475 480

aga aga gta cat gag cag tac tgg ttc ggc tat tcc aag aat gaa gaa 1488  
Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu  
485 490 495

gag tta aga aaa aat gag cta cac cac aag cac tgg tgg aat cca atg 1536  
Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met  
500 505 510

gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac 1584  
Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr  
515 520 525

ggg gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat 1632  
Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp  
530 535 540

gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct 1680  
Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro  
545 550 555 560

gta ttc ctc acc gag ggg gac gga acc gtt ccg ctc gtg gcg cat tca 1728  
Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser  
565 570 575

atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga 1776  
Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly  
580 585 590

att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat 1824  
Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
595 600 605

ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctc ggc agc 1872  
Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
610 615 620

gcg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat 1920  
Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
625 630 635 640

ctc gtc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag 1968  
Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
645 650 655



0-06-1999

10-6-99

EP99111321.8

018 5689334

+49 89 2395

SPEC

19

atg ecc ttc cca atg taa  
Met Pro Phe Pro Met  
660

1986

20

```
<210> 2
<211> 661
<212> PRT
<213> Saccharomyces cerevisiae
<400> 2
```

Met	Gly	Thr	Leu	Phe	Arg	Arg	Asn	Val	Gln	Asn	Gln	Lys	Ser	Asp	Ser
1				5				10						15	
Asp	Glu	Asn	Asn	Lys	Gly	Gly	Ser	Val	His	Asn	Lys	Arg	Glu	Ser	Arg
			20					25					30		
Asn	His	Ile	His	His	Gln	Gln	Gly	Leu	Gly	His	Lys	Arg	Arg	Arg	Gly
		35					40					45			
Ile	Ser	Gly	Ser	Ala	Lys	Arg	Asn	Glu	Arg	Gly	Lys	Asp	Phe	Asp	Arg
	50					55					60				
Lys	Arg	Asp	Gly	Asn	Gly	Arg	Lys	Arg	Trp	Arg	Asp	Ser	Arg	Arg	Leu
65					70					75					80
Ile	Phe	Ile	Leu	Gly	Ala	Phe	Leu	Gly	Val	Leu	Leu	Pro	Phe	Ser	Phe
			85					90					95		
Gly	Ala	Tyr	His	Val	His	Asn	Ser	Asp	Ser	Asp	Leu	Phe	Asp	Asn	Phe
			100					105					110		
Val	Asn	Phe	Asp	Ser	Leu	Lys	Val	Tyr	Leu	Asp	Asp	Trp	Lys	Asp	Val
		115					120					125			
Leu	Pro	Gln	Gly	Ile	Ser	Ser	Phe	Ile	Asp	Asp	Ile	Gln	Ala	Gly	Asn
	130					135					140				
Tyr	Ser	Thr	Ser	Ser	Leu	Asp	Asp	Leu	Ser	Glu	Asn	Phe	Ala	Val	Gly
145					150					155					160
Lys	Gln	Leu	Leu	Arg	Asp	Tyr	Asn	Ile	Glu	Ala	Lys	His	Pro	Val	Val
				165				170					175		
Met	Val	Pro	Gly	Val	Ile	Ser	Thr	Gly	Ile	Glu	Ser	Trp	Gly	Val	Ile
			180					185					190		
Gly	Asp	Asp	Glu	Cys	Asp	Ser	Ser	Ala	His	Phe	Arg	Lys	Arg	Leu	Trp
	195						200					205			
Gly	Ser	Phe	Tyr	Met	Leu	Arg	Thr	Met	Val	Met	Asp	Lys	Val	Cys	Trp
	210					215					220				
Leu	Lys	His	Val	Met	Leu	Asp	Pro	Glu	Thr	Gly	Leu	Asp	Pro	Pro	Asn
225					230					235					240
Phe	Thr	Leu	Arg	Ala	Ala	Gln	Gly	Phe	Glu	Ser	Thr	Asp	Tyr	Phe	Ile
			245					250					255		
Ala	Gly	Tyr	Trp	Ile	Trp	Asn	Lys	Val	Phe	Gln	Asn	Leu	Gly	Val	Ile
			260					265					270		
Gly	Tyr	Glu	Pro	Asn	Lys	Met	Thr	Ser	Ala	Ala	Tyr	Asp	Trp	Arg	Leu
	275						280					285			
Ala	Tyr	Leu	Asp	Leu	Glu	Arg	Arg	Asp	Arg	Tyr	Phe	Thr	Lys	Leu	Lys
	290					295					300				
Glu	Gln	Ile	Glu	Leu	Phe	His	Gln	Leu	Ser	Gly	Glu	Lys	Val	Cys	Leu
305					310					315					320

Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp  
325 330 335  
Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
340 345 350  
Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
355 360 365  
Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
370 375 380  
Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
385 390 395 400  
Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
405 410 415  
Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
420 425 430  
Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
435 440 445  
Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
450 455 460  
Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
465 470 475 480  
Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu  
485 490 495  
Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met  
500 505 510  
Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr  
515 520 525  
Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp  
530 535 540  
Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro  
545 550 555 560  
Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser  
565 570 575  
Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly  
580 585 590  
Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
595 600 605  
Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
610 615 620  
Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
625 630 635 640  
Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
645 650 655  
Met Pro Phe Pro Met  
660

<210> 3  
<211> 2312  
<212> genomic DNA  
<213> Schizosaccharomyces pombe  
<400> 3

ATGGCGTCTT CCAAGAAGAG CAAACTCAT AAGAAAAGA AAGAAGTCAA	50
ATCTCCTATC GACTTACCAA ATTCAAAGAA ACCAACTCGC GCTTTGAGTG	100
AGCAACCTTC AGCGTCCGAA ACACAATCTG TTTCAAATAA ATCAAGAAAA	150
TCTAAATTTG GAAAAAGATT GAATTTTATA TTGGGCGCTA TTTTGGGAAT	200
ATGGCGTGCT TTTTFTTTCG CTGTTGAGCA CGACAATGCT GTTTTCGACC	250
CTGCTACGTT AGATAAATTT GGAATATGC TAGGCTCTTC AGACTTGTTT	300
GATGACATTA AAGGATATTT ATCTTATAAT GTGTTTAAGG ATGCACCTTT	350
TACTACGGAC AAGCCITCGC AGTCTCCTAG CGGAAATGAA GTTCAAGTTG	400
GTCTTGATAT GTACAATGAG GGATATCGAA GTGACCATCC TGTIATTATG	450
GTTCTTGGTG TTATCAGCTC AGGATTAGAA AGTTGGTCGT TTAATAATTG	500
CTCGATTCCT TACTTTAGGA AACGTCTTTG GGGTAGCTGG TCTATGCTGA	550
AGGCAATGTT CCTTGACAAG CAATGCTGGC TTGAACATTT AATGCTTGAT	600
AAAAAAACCG GCTTGGATCC GAAGGGAATT AAGCTGCGAG CAGCTCAGGG	650
GTTTGAAGCA GCTGATTTTT TTATCACGGG CTATTGGATT TGGAGTAAAG	700
TAATTGAAAA CCTTGCTGCA ATTGGTTATG AGCCTAATAA CATGTTAAGT	750
GCTTCTTACG ATTGGCGGTT ATCATATGCA AATTTAGAGG AACGTGATAA	800
ATATTTTTCA AAGTTAAAAA TGTTCAATGA GTACAGCAAC ATTGTACATA	850
AGAAAAAGGT AGTGTTGATT TCTCACTCCA TGGGTTTACA GGTTACGTAC	900
TATTTTTTTA AGTGGGTTGA AGCTGAGGGC TACGGAAATG GTGGACCGAC	950
TTGGGTTAAT GATCATATTG AAGCATTTAT AAATGTGAGT CTCGATGGTT	1000
GTTTGACTAC GTTCTAACT TTTGAATAGA TATCGGGATC TTTGATTGGA	1050
GCACCCAAAA CAGTGGCAGC GCTTTTATCG GGTGAAATGA AAGATACAGG	1100
TATTGTAATT ACATTAAACA TGTTAATATT TAATTTTTGC TAACCGTTTT	1150
AAGCTCAATT GAATCAGTTT TCGGTCTATG GGTAAAGCAAT AAATTGTTGA	1200
GATTTGTTAC TAATTTACTG TTTAGTTTGG AAAAATTTTT TTCCCGTTCT	1250
GAGGTATATT CAAAAATACA AATGTGCTCT ACTTTTTCTA ACTTTTAATA	1300
GAGAGCCATG ATGGTTCGCA CTATGGGAGG AGTTAGTTCT ATGCTTCCTA	1350
AAGCAGGCCA TGTGTATGG GGAAATGCCA GTTGGGTAAG AAATATGTGC	1400

TGTTAATTTT	TTATTAATAT	TTAGGCTCCA	GATGATCTTA	ATCAAACAAA	1450
TTTTCCTCAAT	GGTGCAATTA	TTGATATAG	AGAAGACATT	GATAAGGACC	1500
ACGATGAATT	TGACATAGAT	GATGCATTAC	AATTTTAAA	AAATGTTACA	1550
GATGACGATT	TTAAAGTCAT	GCTAGCGAAA	AATTATTCCC	ACGGTCTTGC	1600
TTGGACTGAA	AAAGAAGTGT	TAAAAAATAA	CGAAATGCCG	TCTAAATGGA	1650
TAAATCCGCT	AGAAGTAAGA	ACATTAAAGT	TACTAAATTA	TACTAACCCA	1700
AATAGACTAG	TCTTCCTTAT	GCTCCTGATA	TGAAAATTTA	TTGCGTTCAC	1750
GGGGTCGGAA	AACCAACTGA	GAGAGGTTAT	TATTATACTA	ATAATCCTGA	1800
GGGGCAACCT	STCATTGATT	CCTCGGTTAA	TGATGGAACA	AAAGTTGAAA	1850
ATGTGAGAGA	ATTTATGTTT	CAAACATTCT	ATTAAGTGT	TTATTAGGGT	1900
ATTGTTAIGG	ATGATGGTGA	TGGAACCTTA	CCAATATTAG	CCCTTGTTTT	1950
GGTGTGCAAT	AAAGTTTGGC	AAACAAAAG	GTTTAATCCT	GCTAATACAA	2000
GTATCACAAA	TTATGAAPTC	AAGCATGAAC	CTGCTGCGTT	TGATCTGAGA	2050
GGAGGACCTC	GCTCGGCAGA	ACACGTCGAT	ATACTTGGAC	ATTCAGAGCT	2100
AAATGTATGT	TCATTCTACC	TTACAAATTT	CTATTACTAA	CTCTTGAAAT	2150
AAGGAAATTA	TTTTAAAAGT	TTCATCAGGC	CATGGTGAAT	CGGTACCAA	2200
CCGTATATA	TCAGATATCC	AGTACGGACA	TAAGTTTTGT	AGATTGCAAT	2250
TAACTAACTA	ACCGAACAGG	GAAATAATAA	ATGAGATAAA	TCTCGATAAA	2300
CCTAGAAATT	AA				2312

<210> 4  
<211> 3685  
<212> genomic DNA  
<213> Arabidopsis thaliana  
<400> 4

ATGCCCTTA TTCATCGGAA AAAGCCGACG GAGAAACCAT CGACGCCGCC	50
ATCTGAAGAG GTGGTGCACG ATGAGGATTC GCAAAAGAAA CCACACGAAT	100
CTTCCAAATC CCACCATAAG AAATCGAACG GAGGAGGGAA GTGGTCGTGC	150
ATCGATTCTT GTTGTGGTGT CATTTGGGTGT GTGTGTGTAA CCTGGTGGTT	200
TCTTCTCTTC CTTTACAACG CAATGCCTGC GAGCTTCCCT CAGTATGTAA	250
CGGAGCGAAT CACGGGTCCCT TTGCCTGACC CGCCCGGTGT TAAGCTCAAA	300
AAAGAAGGTC TTAAGGCGAA ACATCCTGTT GTCTTCATTC CTGGGATTGT	350
CACCGGTGGG CTCGAGCTTT GGGAAGGCAA ACAATGCGCT GATGGTTTAT	400
TTAGAAAACG TTTGTGGGGT GGAACTTTTC GTGAAGTCTA CAAAAGGTGA	450
GCTCAACAAT TCTCACTCTT CCTTTATATT GGGATTGGA TTGCATCTGA	500
TGAGATCACG CACTTGTTCG TTCTTCAACA TCACTCAAAC TTTAATTCCA	550
TGTTTGCTCG TCTTACTCTT TACTTTTTTT TTTTTTTGAT GTGAAACGCT	600
ATTTTCTTAA GAGACTATTT CTGTATGTGT AAGGTAAGCG TTCCAAGGAC	650
GTAATTGGCT TGGACTATTT CTGTTTGATT GTTAACTTTA GGATATAAAA	700
TAGCTGCCTT GGAATTTCAA GTCATCTTAT TGCCAAATCT GTTGCTAGAC	750
ATGCCCTAGA GTCCGTTTCAT AACAAAGTTAC TTCCTTTACT GTCGTTGCGT	800
GTAGATTTAG CTTTGTGTAG CGTATAATGA AGTAGTGTTC TATGTTTTGT	850
TGGGAATAGA GAAGTTCTAA CTACATCTGT GGAAAGTGTG TTCAGGCTGT	900
GATAGAGGAC TGTTGCTTTA TTATTCAACT ATGTATATGT GTAATTAAAG	950
CTAGTTCCTT TTTGATCTTT CAGCTCAATG TGCTTTTCTC AATTTTTTTC	1000
TCAATTTCAA AGTTTCACAT CGAGTTTATT CACATGTCTT GAATTTCTGC	1050
CATCCTCGTT CTGTTATCCA GCTTTGAACT CCTCCCGACC CTGCTATGGA	1100
TATATTAAAA AAAAAGTGTT TTGTGGGTTG CATCTTTGTT ACGATCTGCA	1150
TCTTCTTCTT TCGGCTCAGT GTTCATGTTT TTGCTATGGT AGAGATGGGC	1200
AATGTTATTG TTGATGGTAA CAGTGGTATA GTTGATAGTA TCTTAACTAA	1250
TCAATTATCT CTTTGATTCA GGCCTCTATG TTGGGTGGAA CACATGTCAC	1300
TTGACAATGA AACTGGGTTG GATCCAGCTG GTATTAGAGT TCGAGCTGTA	1350

TCAGGACTCG	TGGCTGCTGA	CTACTTTGCT	CCTGGCTACT	TTGTCTGGGC	1400
AGTGCTGATT	GCTAACCTTC	CACATATTGG	ATATGAAGAG	AAAAATATGT	1450
ACATGGCTGC	ATATGACTGG	CGGCTTTCGT	TTCAGAACAC	AGAGGTTCTT	1500
TTCTCATCGT	TCTTTCTATT	ATTCTGTTCC	ATGTTACGTT	TCTTTCTTCA	1550
TTACTTAAGG	CITAAATATG	TTTCATGTTG	AATTAATAGG	TACGTGATCA	1600
GACTCTTAGC	CGTATGAAAA	GTAATATAGA	GTTGATGGTT	TCTACCAACG	1650
GTGGAAAAAA	AGCAGTTATA	GTTCCGCATT	CCATGGGGGT	CTTGTATTTT	1700
CTACATTTTA	TGAAGTGGGT	TGAGGCACCA	GCTCCTCTGG	GTGGCGGGGG	1750
TGGGCCAGAT	TGGTGTGCAA	AGTATATTAA	GGCGGTGATG	AACATTGGTG	1800
GACCATTCT	TGGTGTGCAA	AAAGCTGTTG	CAGGGCTTTT	CTCTGCTGAA	1850
GCAAAGGATG	TTGCAGTTGC	CAGGTATTGA	ATATCTGCTT	ATACTTTTGA	1900
TGATCAGAAC	CTTGGCTCTG	GAAC TCAAAG	TTATTCTACT	AAATATCAAT	1950
TCTAATAACA	TIGCTATATT	ATCGCTGCAA	CTGACATTGG	TTGATTATTT	2000
TTGCTGCTTA	TGTAAGTGAA	ACTCTCTTGA	GATTAGACAA	ATGATGAATT	2050
GATAATTCTT	ACGCATTGCT	CTGTGATGAC	CAGTTTCTTA	GCTTCGACGA	2100
TAAACATTGT	CATACTGTCT	TTTGGAGGGC	ATTGAATTTT	GCTATGGAAA	2150
GCGCTGGAGC	TTCCATGCTT	GCATTCTTTA	CCAATTAGCG	TTATTCTGCT	2200
TCTTTCAATT	TTCTTGATA	TGCATCTATG	GTCTTTTATT	TCTTCTTAAT	2250
TAAAGACTCG	TTGGATTAGT	TGCTCTATTA	GTCACTTGGT	TCCTTAATAT	2300
AGAACTTTAC	TTTCTTCGAA	AATTGCAGAG	CGATTGCCCC	AGGATTCTTA	2350
GACACCGATA	TATTTAGACT	TCAGACCTTG	CAGCATGTAA	TGAGAATGAC	2400
ACGCACATGG	GACTCAACAA	TGTCTATGTT	ACCGAAGGGA	GGTGACACGA	2450
TATGGGGCGG	GCTTGATTGG	TCACCGGAGA	AAGGCCACAC	CTGTTGTGGG	2500
AAAAAGCAAA	AGAACAACGA	AACTTGTGGT	GAAGCAGGTG	AAAACGGAGT	2550
TTCCAAGAAA	AGTCCTGTTA	ACTATGGAAG	GATGATATCT	TTTGGGAAAG	2600

CTATAGATCT ACTACATTAT GTTGCTCCTA AGATGATGGC GCGTGGTGCC	2900
GCTCATTCTT CTTATGGAAT TGCTGATGAT TTGGATGACA CCAAGTATCA	2950
AGATCCCAAA TACTGGTCAA ATCCGTTAGA GACAAAGTAA GTGATTTCTT	3000
GATTCCAACCT GTATCCTTCG TCCTGATGCA TTATCAGTCT TTTTGTTTTT	3050
GGTCTTGTTG GATATGGTTT TCAGCTCAAA GCCTTACAAAG CTGTTTCTGA	3100
GCCTTTCTCA AAAAGGCTTG CTCAGTAATA TTGAGGTGCT AAAGTTGATA	3150
CATGTGACTC TTGCTTATAA ATCCTCCGTT TGGTTTGTTT TGCTTTTTC	3200
GATTACCGAA TGCTCCTGAG ATGGAAATCT ACTCATTATA CGGAGTGGGG	3250
ATACCAACGG AACGAGCATA CGTATACAAG CTTAACCAGT CTCCCGACAG	3300
TTGCATCCCC TTTCAGATAT TCACTTCTGC TCACGAGGAG GACGAAGATA	3350
GCTGTCTGAA AGCAGGAGTT TACAATGTGG ATGGGGATGA AACAGTACCC	3400
GTCCTAAGTG CCGGGTACAT GTGTGCAAAA GCGTGGCGTG GCAAGACAAG	3450
ATTCAACCCT TCCGGAATCA AGACTTATAT AGAGAATAC AATCACTCTC	3500
CGCCGGCTAA CCTGTTGGAA GGGCGCGGGA CGCAGAGTGG TGCCCATGTT	3550
GATATCATGG GAAACTTTGC TTTGATCGAA GATATCATGA GGGTTGCCGC	3600
CGGAGGTAAC GGGTCTGATA TAGGACATGA CCAGGTCCAC TCTGGCATAT	3650
TTGAATGGTC GGAGCGTATT GACCTGAAGC TGTGA	3685



0-06-1999

-MUNICHEN 03  
00 1110 10:41

110- 6-99

LUDWIG DR

EP99111321.8

018 568939

1110 000000

+49 89 23

SPEC

27

<210> 5  
<211> 402  
<212> cDNA  
<213> Arabidopsis thaliana  
<220>  
<221> CDS  
<222> (120)..(402)  
<400> 5

AGAAACAGCTCTTTGTCTCT CTCGACTGATCTAACAATCC CTAATCTGTGTTCTAAATTC 60  
CTGGACGAGATTTGACAAAG TCCGTATAGCTTAACCTGGT TTAATTTCAAGTGACAGAT 119  
ATG CCC CTT ATT CAT CGG AAA AAG CCG ACG GAG AAA CCA TCG ACG CCG 167  
Met Pro Leu Ile His Arg Lys Lys Pro Thr Glu Lys Pro Ser Thr Pro  
CCA TCT GAA GAG GTG GTG CAC GAT GAG GAT TCG CAA AAG AAA CCA CAC 215  
Pro Ser Glu Glu Val Val His Asp Glu Asp Ser Gln Lys Lys Pro His  
GAA TCT TCC AAA TCC CAC CAT AAG NAA TCG AAC GGA GGA GGG AAG TGG 263  
Glu Ser Ser Lys Ser His His Lys ??? Ser Asn Gly Gly Gly Lys Trp  
TCG TGC ATC GAT TCT TGT TGT TGG TTC ATT GGG TGT GTG TGT GTA ACC 311  
Ser Cys Ile Asp Ser Cys Cys Trp Phe Ile Gly Cys Val Cys Val Thr  
TGG TGG TTT CTT CTC TTC CTT TAC AAC GCA ATG CCT GCG AGC TTC CCT 359  
Trp Trp Phe Leu Leu Phe Leu Tyr Asn Ala Met Pro Ala Ser Phe Pro  
CAG TAT GTA ACG GAG CCG AAT CAC GNG TCC TTT GCC TTA CCC G 402  
Gln Tyr Val Thr Glu Pro Asn His ??? Ser Phe Ala Leu Pro

<210> 6  
<211> 643  
<212> cDNA  
<213> ZPA MAYA  
<220>  
<221> CDS  
<222> (1)..(402)  
<400> 6

```
CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 46
Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
   1                               5               10               15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA 96
Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
                   20                   25                   30

TGG CGT GGC AAA ACT CGT TTC AGC CCT GCC GGC AGC AAG ACT TAC GTG 144
Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
                   35                   40                   45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC 192
Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
   50                               55               60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT 240
Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
  65                               70               75               80

GAG GAC GTC ATC AGA ATA GCT GCT GGG GCA ACC GGT GAG GAA ATT GGT 288
Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
                   85                   90                   95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA 336
Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
                   100                  105                  110

TTG AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTCGCTTCC 391
Leu Lys Leu
   115

AAAGTGTCTCT GCCTGAGTGC AACTCTGGAT TTTGCTTAAA TATTGTAATT TTTCACGC 449

TTCATTCGTC CCTTTGTCAA ATTTACATTT GACAGGACGC CAATGCGATA CGATGTTG 507

TACCGCTATT TTCAGCATTG TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC 565

TGAAATTGTG TAGTCGTTTT CTTTACGATT TAATANCAAG TGGCGGAGCA GTGCCCCA 623

AGCNAAAAAA AAAAAAAAAA 643
```

<210> 7  
 <211> 115  
 <212> PRT  
 <213> Zea mays  
 <400> 7

Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly  
 1 5 10 15

Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly  
 20 25 30

Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val  
 35 40 45

Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly  
 50 55 60

Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile  
 65 70 75 80

Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly  
 85 90 95

Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys  
 100 105 110

Leu Lys Leu  
 115

10-06-1999

EP991113218

015 8939

+49 89 23

SPEC

30

<210> 8  
<211> 616  
<212> cDNA  
<213> *Neurospora crassa*  
<400> 8

GGTGGCGAAG ACGANGGCGG AAGTTGGAGG CTAACGAGAA TGACNCTCGG	50
AGATGGATCT ACCCTCTAGA GACACGACTA CCNTTGCACC CAGCCTCAAG	100
GTNTACNGTT TNTATGGGTA GGAAGCCGAC GGAGCGAGCC TACATCTATC	150
TGCGGCCCCGA TCCCGGGACG ACAACGCATC TTTAGATGAC GATCGATACG	200
ACTTTGACTN AGGGGCACAT TGACCACGGT GTGATTTTGG GCGAAGGCGA	250
TGGCACAGTG AACCTTATGA GTTTGGGGTA CCTGTGCAAT AAGGGGTGGA	300
AAATGAAGAG ATACAATCCT GCGGGCTCAA AAATAACCGT GGTCGAGATG	350
CCGCATGAAC CAGAACGGT CAATCCGAGA GGAGGGCCGA ATACGGCGGA	500
CTTAAATATG TAGAAAAGGT TGAAATTTAT GAAGAGTAAT TAAATACGGC	550
ACATAGGTTA CTCAATAGTA TGACTAATTA AAAAAAATT TTTTTTCTAA	600
AAAAAAAAAA AAAAAA	616

<210> 9  
 <211> 1562  
 <212> genomic DNA  
 <213> Arabidopsis thaliana  
 <400> 9

ATGAAAAAA TATCTTCACA TTATTCGGTA GTCATAGCGA TACTCGTGTG	50
GGTGACGATG ACCTCGATGT GTCAAGCTGT GGGTAGCAAC GTGTACCCTT	100
TGATTCTGGT TCCAGGAAAC GGAGSTAACC AGCTAGAGGT ACGGCTGGAC	150
AGAGAATACA AGCCAAGTAG TGTCTGGTGT AGCAGCTGGT TATATCCGAT	200
TCATAAGAAG AGTGGTGGAT GGTTTAGGCT ATGGTTCGAT GCAGCAGTGT	250
TATTGTCTCC CTTCAACAGG TGCTTCAGCG ATCGAATGAT GTTGTACTAT	300
GACCCTGATT TGGATGATTA CCAAAATGCT CCTGGTGTCC AAACCCGGGT	350
TCCTCATTTT GGTTCGACCA AATCACTTCT ATACCTCGAC CCTCGTCTCC	400
GGTLAGTACT TTCCAAGATA TATCATTTTG GGACATTTGC ATAATGAACA	450
AAATAGACAT AAATTTGGGG GATTATTGTT ATATCAATAT CCATTTATAT	500
GCTAGTCGGT AATGTGAGTG TTATGTTAGT ATAGTTAATG TGAGTGTTAT	550
GTGATTTTCC ATTTTAAATG AAGCTAGAAA GTTGTCGTTT AATAATGTTG	600
CTATGTCATG AGAATTATAA GGACACTATG TAAATGTAGC TTAATAATAA	650
GGTTTGATTT GCAGAGATGC CACATCTTAC ATGGAACATT TGGTGAAAGC	700
TCTAGAGAAA AAATGCGGGT ATGTTAACGA CCAAACCATC CTAGGAGCTC	750
CATATGATTT CAGGTACGGC CTGGCTGCTT CGGGCCACCC GTCCCGTGTA	800
GCCTCACAGT TCCTACAAGA CCTCAAACAA TTGGTGGAAG AAACCTAGCAG	850
CGAGAACGAA GGAAAGCCAG TGATACTCCT CTCCCATAGC CTAGGAGGAC	900
TTTCGTCTCT CCATTTCTCT AACCGTACCA CCCCTTCATG GCGCCGCAAG	950
TACATCAAAC ACTTTGTTGC ACTCGCTGCG CCATGGGGTG GGACGATCTC	1000
TCAGATGAAG ACATTTGCTT CTGGCAACAC ACTCGGTGTC CCTTTAGTTA	1050
ACCCTTTGCT GGTCAGACGG CATCAGAGGA CCTCCGAGAG TAACCAATGG	1100
CTACTTCCAT CTACCAAAGT GTTTCACGAC AGAACTAAAC CGCTTGTCGT	1150
AACTCCCCAG GTTAACTACA CAGCTTACGA GATGGATCGG TTTTTTGCAG	1200
ACATTGGATT CTCACAAGGA GTTGTGCCTT ACAAGACAAG AGTGTTGCCT	1250

TTAACAGAGG AGCTGATGAC TCCGGGAGTG CCAGTCACTT GCATATATGG	1300
GAGAGGAGTT GATACACCGG AGGTTTTGAT GTATGGAAAA GGAGGATTCG	1350
ATAAGCAACC AGAGATTAAG TATGGAGATG GAGATGGGAC GGTTAATTTG	1400
GCGAGCTTAG CAGCTTTGAA AGTCGATAGC TTGAACACCG TAGAGATTGA	1450
TGGAGTTTCG CATACTCTA TACTTAAAGA CGAGATCGCA CTTAAAGAGA	1500
TTATGAAGCA GATTTCAATT ATTAATTATG AATTAGCCAA TGTTAATGCC	1550
GTCAATGAAT GA	1562

<210> 10  
<211> 3896  
<212> genomic DNA  
<213> Arabidopsis thaliana  
<400> 10

ATGGGAGCGA	ATTGGAATC	AGTAACGGCT	TCCTTCACCG	TCATCGCCGT	50
TTTTTCTTG	ATTTCGGTG	GCCGAACGTC	GGTGGAGGAT	GAGACCGAGT	100
TTCACGCCGA	CTACTCGAAG	CTATCGGGTA	TAATCATTCC	GGGATTTGCG	150
TCGACGCAGC	TACGAGCGTG	GTCGATCCTT	GACTGTCCAT	ACACTCCGT	200
CGACTTCAAT	CCGCTCGACC	TCGTATGGCT	AGACACCACT	AAGGTCCGTG	250
ATCTTCATTT	CCTTCGCTCC	TTATTCTGTC	GGTCGAGTCA	CTTGTTGATG	300
AATTCCAAGC	GAAATATAGC	AATGAAGCAT	GTCTCGTCTC	TCTTATTGAT	350
TCGTTCATTA	GTCAACAGTG	ACGCTTCTGA	ATCTGAGTTT	AGAGTCATAT	400
AAAACAGCTG	ACTCGGCGAG	TGTTTCCCAT	CGCTTTTGGT	TCGCTAAATG	450
TAGCGCAATG	AATGTGTAAT	TAGTCTGCGC	TTTTTATTCA	ACTAGATCTG	500
CAAGTTTTTC	AGAGTGCTCA	ATAGTAGTTA	GAAATGTTA	GGTCATTTTA	550
CTTGTCATTT	GTGATTCTTT	TGGTTGTTGC	TTACTGATCG	ACGTGATGGA	600
TGGTTTACAG	CTTCTTTCTG	CTGTCAACTG	CTGGTTTAAAG	TGTATGGTGC	650
TAGATCCTTA	TAATCAAACA	GACCATCCCG	AGTGTAAAGTC	ACGGCCTGAC	700
AGTGGTCTTT	CAGCCATCAC	AGAATTGGAT	CCAGGTTACA	TAACAGGTA	750
TTTCGGATTT	TTCTTTCTTT	TGAGTTTTCT	TCAATTTGAT	ATCATCTTGT	800
TGTGATATAA	TATGGCTAAG	TTCAATTAAT	GGTCAATTT	TCAGGTCCTC	850
TTTCTACTGT	CTGAAAGAG	TGGCTTAAGT	GGTGTGTTGA	GTTTGGTATA	900
GAAGCAAATG	CAATTGTGCG	TGTTCCATAC	GATTGGAGAT	TGTCACCAAC	950
CAAAATGGAA	GAGCGTGACC	TTTACTTTCA	CAAGCTCAAG	TTAGTCCTTA	1000
TCAGGCTAAT	GTCTTTTATC	TTCTCTTTTT	ATGTAAGATA	AGCTAAGAGC	1050
TCTGGTCTGC	TTCTTTTTTG	CAGGTGAGCC	TTTGAAACTG	CTTTAAAACT	1100
CCGTGGCGGC	CCTTCTATAG	TATTTGCCCA	TTCAATGGGT	AATAATGTCT	1150
TCAGATACTT	TCTGGAATGG	CTGAGGCTAG	AAATTGCACC	AAAACATTAT	1200
TTGAAGTGGC	TTGATCAGCA	TATCCATGCT	TATTTGCTG	TTGGTACCGG	1250
CCTACTATCC	TTAAGTTACC	ATTTTATTTT	TTCTCTAATT	GGGGGAGTTA	1300
TGTTGTGACT	TACTGGATTG	AGCTCGATAC	CTGATTTGTT	GTTGATTAG	1350
GAGCTCCTCT	TCTTGGTTCT	GTTGAGGCAA	TCAAATCTAC	TCTCTCTGGT	1400
GTAACGTTTG	GCCTTCCTGT	TTCTGAGGTG	ACCTCTGACT	TCTCTTTAGT	1450
TTTAAGTAGT	TGATATCAAC	CAGGTCTTAT	AACTCACTGG	ATTTTCCTTT	1500
TGAAAGTATT	ACTTTTGTTA	ATTGAAGTGC	TGTACGCGAT	ATGGTATCTG	1550
TAGATCTTGA	AGTGCTAGTT	ATCAAAGAAC	ATATTGTGGG	TAGTATACCT	1600
GTCAAGCGCC	TTAGCTAATA	CAACCAAACC	ACATGTACAC	TGATTTAOTT	1650
TTCAAGATTAT	TATGGTAGAC	TTTAAGTTGA	GAAGAAACTT	TGACTGAAAT	1700
CTTTTATTTT	TAATAGGCTA	TGATTTGTTT	ATTGAAATCA	TGTGACATAT	1750
TGACATGCGC	TTCTCATGTT	TTTTGTTGGC	AAGGCTTCAG	GGAAGTGCTC	1800
GGTTGTTGTC	CAATTCTTTT	GCGTCGTCT	TGTGGCTTAT	GCCATTTTCA	1850
AAGAATTGCA	AGGGTGATAA	CACATTCTGG	ACGCATTTT	CTGGGGGTGC	1900
TGCAAGAAA	GATAAGCGCG	TATACCACTG	TGATGAAGAG	GAATATCAAT	1950
CAAAATATTC	TGGCTGGCCG	ACAAATATTA	TTAACATTGA	AATTCCTTCC	2000
ACTAGCGGTT	AGACTCTGTA	TATGCAACTG	TAACACTAAC	AAAAGTTTCA	2050
CCAAGAATGT	TCACTCTCAT	ATTTGCTTCC	TTTGATGTGT	ATCCATCAGT	2100
TACAGAAACA	GCTCTAGTCA	ACATGACCAG	CATGGAATGT	GGCCTTCCCA	2150
CCCTTTTGTC	TTTCACAGCC	CGTGAAGTAG	CAGATGGGAC	TCCTTTTCAA	2200
GCAATAGAAG	ACTATGACCC	AGATAGCAAG	AGGATGTTAC	ACCAGTTAAA	2250
GAAGTACGTA	CCTTTCTTTG	TGATAAGAAA	TATTGCTCAT	CGATCATCAC	2300
TTGCTGGCTT	CTTGTAAGTC	AAATTGTTTT	GTTTAAATCT	CTATATCAAT	2350
TGTTTCATATG	CTTTGTCTTT	CTTACTATAA	GAAACAAGTA	TAATCAGAAA	2400
CCTTATATTT	GATTATCAGT	TCTCTCCTTA	TATTATGGAA	TGTCTTTTTC	2450
GTTTACAGTT	ATGAATGCAA	AAGGGGTAT	TTTAGTTGAT	TGATTCTCTC	2500
ATCTCTAGT	TTGTTTTGAC	TAATAGCGTC	AATTTTGTTT	TTCTAGCAAA	2550
TCTTTGTGAA	TTATATATAA	CATGCTAACT	ATACTTTTCA	GGTGTATCA	2600
TGATGACCTT	GTTTTTAATC	CTCTGACTCC	TTGGGAGAGA	CCACCTATAA	2650
AAAATGTATT	TTGCATATAT	GGTGCTCATC	TAAAGACAGA	GGTATGATGC	2700
ATTCTCAATA	TCACATTATG	CGTTGACTTT	GTTATTATAT	TCCCCATTG	2750

10-06-1999 MUENCHEN 03  
09 1110 10:40

6-99 :  
DNR EP99111321.8

018 939  
000000

+49 89 238

SPEC

34

GTTTGCAATA	TCTTTTTGAA	TTATGATTTA	TCTTCTCCCT	TGCATCTTAT	2800
GCTATTAAGC	GTTAAAGGTA	CTAAATGTAT	GAAGCTGTCT	GTCATAGGTT	2850
GGTTATTACT	TTGCCCCAAG	TGGCAAACTT	TATCTTGATA	ATGGGATCAT	2900
CACGGATATC	ATTTATGAAA	CTGAAGGTTT	CCTCGTGTCA	AGGTAATTTT	2950
CCGCAATGGC	AGAAGTAAAA	CAGGAAGGCA	AAGTCTTCTG	TATCAGTCTA	3000
G'GGCATGTT	ATCTCAGTTG	CATAAGCAAA	TTATTAACA	ACTAAAATTT	3050
AAGTACTTTT	TTATCATTCC	TTTTGAGCTT	AGTGGATGAT	CAGTGGCTTA	3100
AACTGGGAAG	AGGTGTTGCA	TGAAACATGA	CACTTGTATC	AAAGATAACT	3150
AGCAAAACAA	AACTAACCCA	TTTCTGAATT	TCATATTATT	AGGAGTAGTC	3200
GTGCTTTTAA	AAAATTTGTT	TTAAGAAACC	GAAAAACTAG	TTCATATCTT	3250
GATTGTGCAA	TATCTGCAGG	TCTGGAACCT	TGGTTGATGG	GAACGCTGGA	3300
CCTATAACTG	GGGATGAGAC	GGTAAGCTCA	GAAGTTGOTT	TTGAAATTAT	3350
CTTCTTGCAA	ACTACTGAAG	ACTAAGATAA	TACTTGCTTC	TGGAACACTG	3400
CTTGCTATGT	TCTCTAGTAC	ACTGCAATAT	TGACTCTCCG	CTACTTTTAT	3450
TGATTATGAA	ATTGATCTCT	TATAGGTACC	CTATCATTCA	CTCTCTTGGT	3500
OCAAGAATTG	GCTCGGACCT	AAAGTTAACA	TAACAATGGC	TCCCAGGTA	3550
CTCTTTTTTA	GTTCCCTCAC	TTATATAGAT	CAAACTTTAA	GTGTACTTTT	3600
CTGGTTATGT	GTTGATTTAC	CTCCAATTTG	TTCTTTCTAA	AAATCATATA	3650
TCTCTGTACT	CCTCAAGAAC	TTGTATTAAT	CTAAACGAGA	TTCTCATTGG	3700
GAAAAATAAA	CAACAGCCAG	AACACGATGG	AAGCGACGTA	CATGTGGAAC	3750
TAAATGTTGA	TCATGAGCAT	GGGTCAGACA	TCATAGCTAA	CATGACAAAA	3800
GCACCAAGGG	TTAAGTACAT	AACCTTTTAT	GAAGACTCTG	AGAGCATTCC	3850
GGGGAAGAGA	ACCCAGTCT	GGGAGCTTGA	TAAAAGTGGG	TATTAA	3896



<210> 11  
<211> 709  
<212> cDNA  
<213> tomato  
<400> 11

CTGGGGCCAA AAGTGAACAT AACAAAGGACA CCACAGTCAG AGCATGATGT	50
TCAGATGTAC AAGTGCATCT AAATATAGAG CATCAACATG GTGAAGATAT	100
CATTCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG	150
AGGATTCTGA AAGTTTTCCTA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT	200
AAAGCAAATC ACAGGAACAT TGTCAGATCT CCAGCTTTGA TGCGGGAGCT	250
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAAG TCCAAGTTTG	300
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AAATGTTTGA	350
GTTGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC	400
TGAGGGATGA GGA CTGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATCG	450
GCTGAACATT GTGAATACAG GTTAGAATAT TCAAATTATA TTTTGCAAAA	500
TATTCTCTTT TTGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA	550
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT	600
TCTACATCTC ACATCCGTGC AACTATGTG TGATATTTAA GAAACTTTGT	650
TTGGCGGAAC AACAAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA	700
TCAGAGAG	709

Fig 1

